



Strokes can lead to seizures.

A person who has a **stroke** and begins having seizures months or even years later may develop epilepsy.

Nearly 10% of stroke survivors suffer from post-stroke seizures.

43% of the time, those seizures happen within 24 hours of the stroke.

Stroke causes nearly 45% of new epilepsy cases for adults 60 and over

Know warning signs for strokes



Learn Seizure First Aid





STAY with the person until they are awake and alert after the seizure.





Keep the person SAFE.

✓ Move or guide away from harm



Turn the person onto their SIDE if they are not awake or aware.

- ✓ Keep airway clear
- ✓ Loosen tight clothes around neck
- ✓ Put something small and soft under the head

Call **911** if...

- Seizure lasts longer than 5 minutes
- Person does not return to their usual
- Person is injured, pregnant, or sick
- Repeated seizures First time seizure
- Difficulty breathing
- Seizure occurs in water





The Epilepsy Foundation and American Stroke Association are working together to raise awareness about the relationship between epilepsy and stroke. Think about making lifestyle changes that can reduce your risk of having a stroke. The #StrokesToSeizures campaign is supported by the CDC/US Dept. of HHS as part of a financial assistance award totaling \$30,000 with 100% funded by the CDC/HHS. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by, CDC/HHS, or the US Government.